


OVAIO

Anni 2005-2017. Sintesi dei dati di incidenza. OVAIO

	FEMMINE
Numero casi	655
% sul totale	2,9
Tasso grezzo	17
Tasso standardizzato	15,6
Rischio cumulativo	1
Età media	63,7
Età mediana	66
Oss/Att AIRTUM (2008-2013)	1,02
Oss/Att Centro (2008-2013)	1,01
Oss/Att Nord (2008-2013)	0,98
Oss/Att Sud (2008-2013)	1,09

Anni 2005-2017. Trend dei dati di incidenza. OVAIO

	FEMMINE		
	Num_F	TSD_F	ES_TSD_F
2005	46	14,85	2,22
2006	53	16,67	2,31
2007	37	11,99	1,99
2008	59	18,46	2,43
2009	50	15,59	2,23
2010	45	14,03	2,12
2011	51	16,00	2,28
2012	48	15,60	2,28
2013	55	16,44	2,25
2014	59	17,65	2,34
2015	49	14,60	2,13
2016	51	15,38	2,20
2017	52	15,48	2,19




Anni 2005-2017. Tassi età specifici di incidenza OVAIO

ETA'	FEMMINE
0-04	0,00
05-09	0,00
10-14	1,67
15-19	1,00
20-24	2,71
25-29	2,13
30-34	6,04
35-39	4,87
40-44	11,55
45-49	23,05
50-54	22,20
55-59	23,72
60-64	20,69
65-69	34,50
70-74	39,40
75-79	47,75
80-84	32,63
85+	43,68




Anni 2005-2017. Sopravvivenza. OVAIO

	2005-09	2010-14	stima 2015-17
1a_F	70,72	69,76	82,97
2a_F	56,55	56,43	65,78
3a_F	48,52	48,83	60,25
4a_F	43,82	43,88	52,00
5a_F	38,52	40,42	45,50



Anno 2017. Prevalenza osservata. OVAIO


Anni	FEMMINE	
	Casi	Tassi
2009	82	28,26
2010	112	38,60
2011	136	46,87
2012	153	52,72
2013	174	59,96
2014	194	66,85
2015	203	69,95
2016	215	74,09
2017	230	79,26



Anni 2005-2017. Frequenza morfologie confermate microscopicamente. OVAIO

FEMMINE	N	%
Sieroso	171	44,42
Mucinoso	40	10,39
Endometrioide	31	8,05
Altri adenocarcinomi	24	6,23
Altre morfologie	32	8,31
Non specificato	87	22,60

Anni 2005-2017. Trend delle morfologie.



FEMMINE

